

État des connaissances sur le Grand tétras (*Tetrao urogallus major*)

Bibliographie actualisée classée par thématiques

(E. Ménoni, 2023)

Statut des populations de Grand Tétras

Adamic, M. (1990). The Capercaillie (*Tetrao urogallus L.*) in Slovenia. In Slovenian with english summary. *Divjipetelin (Tetrao urogallus L.) v Sloveniji. Gozdarski vestnik* 148(2): 67-84.

Artuso, I. and P. Franceschi De (1988). « Il gallo cedrone (*Tetrao urogallus*) in alcuni ambienti forestali dell'alto adige. Osservazioni sugli habitat preferenziali nel periodo della riproduzione e dello sviluppo. » *Bulletin mus. civ. st. nat. verona* 14 : 381-396.

Baines, D. (1990). Capercaillie in Scotland. *Game conservancy annual review* 22 : 153-156.

Ballesteros, F. and L. Robles (2006). *Manual de Conservation y Manejo del Habitat del Urogallo Cantabrico. Primera reimpression.* Madrid.

Bañuelos, M. J., M. Quevedo, and J. R. Obeso. (2004). Conservation of the cantabrian Capercaillie (*Tetrao urogallus cantabricus*) : A challenge at the edge of the species distribution. *Grouse news* 28 : 20-23.

Boakes, E. (2007). Geographic range change in Galliformes. *Grouse news* 33 : 17-18.

Borchtchevski V., Desbrosses R. et Leclercq B. (1992). Les tétraonidés de la taïga d'Arkangelsk (U.R.S.S.). *Bull. Mens. Off. Nat. Chasse*, 166 : 1327.

Borgo, A. and S. M. Mattedi (2011). Habitat di specie e conservazione del Gallo cedrone in Friuli Venezia Giulia e nel Parco Naturale Dolomiti Friulane. *XVI Convegno Italiano di Ornitologia*.

Botev, N., I. Kolev, and N. Ninov. (1993). Basic morphologic signs, feeding and reproduction of Capercaillie (*Tetrao urogallus L.*) in Bulgaria. Pages 144-147. In *Proceedings of XXI IUGB Congress*. 2 : 144-147.

Catusse, M., C. Novoa, E. Ménoni, J. Poirot, and B. Leclercq. 1992. Statut des populations de grand tétras en France. *Bulletin Mensuel de l'Office National de la Chasse* 171 : 14-19.

De-Franceschi, P. F. 1994. Status, geographical distribution and limiting factors of Capercaillie (*Tetrao urogallus*) in Italy. *Gibier Faune Sauvage- Game and Wildlife* 11 : 161-183.

Duriez, O., and E. Ménoni. 2008. Le Grand Tétras (*Tetrao urogallus*) en France : biologie, écologie et systématique. *Ornithos* 14 : 233-243.

Ewing, S. R., M. A. Eaton, T. F. Poole, M. Davies, and S. Haysom. 2012. The size of the Scottish population of Capercaillie *Tetrao urogallus*: results of the fourth national survey. *Bird Study* 59 : 126-138.

Franceschi, D. 1994. Status, Geographical Distribution and Limiting Factors of Capercaillie (*Tetrao urogallus*) in Italy. *Gibier Faune sauvage, Game and Wildflife*. 11 : 161-183.

Gregersen, F., and H. Gregersen. 2009. Ongoing population decline and range contraction in Norwegian forest grouse. *Ornis Norvegica* 32 : 179-189.

Groupe Tétras Jura, 2020, Le Grand Tétras dans le massif jurassien : statut et répartition, période 2012-2020.

Groupe Tétras Vosges. Synthèses annuelles des suivis de la population de Grand Tétras dans le cadre de l'Arrêté Préfectoral de derogation au derangement intentionnel d'espèce protégée sur le massif des Vosges, 2021 et 2022.

Helle, P., V. Belkin, L. Bjudnik, P. I. Danilov, and A. Jakimov. 2003. Changes in grouse populations in Finland and Russian Karelia during recent decades. In finnish, with english summary: Metsäkanalintukannat Suomessa ja Venäjän Karjalassa. Suomen riista 49: 32-43.

Jacob, G., R. Debrunner, F. Gugerli, B. Schmid, and K. Bollmann. 2009. Field surveys of capercaillie (*Tetrao urogallus*) in the Swiss Alps underestimated local abundance of the species as revealed by genetic analyses of non-invasive samples. Conservation Genetic Published on line DOI 10.1007/s10592-008-9794-8 : 12 pp.

Jahren, T., T. Storaas, T. Willebrand, P. Fossland-Moa, and B.-R. Hagen. 2016. Declining reproductive output in capercaillie and black grouse – 16 countries and 80 years. Animal Biology : 363-400.

Kempf, C., N. Lefranc, J. J. Pfeffer, and F. Villaume. 1974. Le grand tétras dans les Vosges. Alauda 42 : 17-38.

Klaus, S., and H. H. Bergmann. 1994. Distribution, status and limiting factors of Capercaillie (*Tetrao urogallus*) in central Europe, particularly in Germany, including evaluation of reintroductions. Gibier Faune Sauvage- Game and Wildlife 11 : 57-80.

Klaus, S., and H. H. Bergmann. 1994. Restauration plan for Capercaillie (*Tetrao urogallus*) in Germany. Gibier Faune Sauvage- Game and Wildlife 11 : 83-96.

Kortland, K. 2008. Capercaillie conservation in the state forests of Scotland. Grouse news 35 : 7-10.

Kortland, K., and A. Peace. 2013. Capercaillie monitoring on the national forest estate in Scotland: 2002 to 2012. Grouse news 45 : 9-16.

Kvasnes, M. A. J., T. Storaas, H. C. Pedersen, S. Bjørk, and E. B. Nilsen. 2010. Spatial dynamics of Norwegian tetraonid populations. Ecology Research DOI 10.1007/s11284-009-0665-7 25 : 367-374.

Leclercq, B. 2008. Le grand tétras *Tetrao urogallus* dans le Jura français : historique et statut actuel. Ornithos 15 : 256-265.

Leclercq, B., E. Ménoni, and M. Montadert. 2012. Réintroduction du Grand tétras par le Parc National des Cévennes. Bilan en 2012, enseignements tirés de ce projet, perspectives d'avenir. ONCFS-Parc National des Cévennes.

Lefranc, N. 1987. Situation du grand tétras (*Tetrao urogallus*) dans le massif vosgien. Bulletin. Mensuel de l'Office National de la Chasse 112 : 5-18.

Lefranc, N., and F. Preiss. 2008. Le grand tétras *Tetrao urogallus* dans les Vosges : historique et statut actuel. Ornithos 15 : 244-255.

Linden, H. 1981. Estimation of juveniles mortality in the Capercaillie, *Tetrao urogallus*, and the Black grouse, *tetrao tetrix*, from indirect evidence. Finnish Game Researches 39:(35-51).

Linden, H. 1983. Variations in clutch size and egg size of Capercaillie and Black Grouse (in finnish, english summary). Suomen Riista 30 : 44-50.

- Linden, H. 1989. Characteristics of tetraonid cycles in Finland. *Finnish Game Research* 46 : 34-42.
- Linden, H., A. Laurila, and M. Wikman. 1990. Predictability of the capercaillie and black grouse density of the next year - wisdom after the event. *Suomen riista* 36 : 81-88.
- Magnani, Y., M. H. Cruveillé, R. Huboux, and P. Collard. 1991. Entre Rhône et Rhin : grand tétras et gélinotte. Statut territorial et évolution. *Bulletin mensuel de l'Office National de la Chasse* 162 : 9-16.
- Ménoni, E. 1994a. Plan de restauration du grand tétras (*Tetrao urogallus*) en France. *Gibier Faune Sauvage- Game and Wildlife* 11 : 159-202.
- Ménoni, E., 1994b. Statut, évolution et facteurs limitants des populations de grand tétras (*Tetrao urogallus*) : synthèse bibliographique. *Gibier Faune Sauvage- Game and Wildlife*. 11: 597-158.
- Ménoni, E. 1996. Location and size of capercaillie *Tetrao urogallus* leks in relation to territories of
- Ménoni, E. 2006. El urogallo en los Pirineos franceses: situacion actual de la especie, evolucion de los efectivos, factores limitantes y gestion de las poblaciones. Pages 79-84 in B. Life, editor. *El urogallo en Espana, Andorra y pirineos franceses. Situacion actual (2005)*.
- Ménoni, E., C. Calenge, E. Dumont-Dayot, and B. Milhau (2014). Estimation des effectifs de Grand tétras dans les Pyrénées françaises. *Gibier Faune sauvage* 304 : 32-36.
- Ménoni, E., M. Catusse, C. Novoa, M. Levet, J. F. Brenot, and P. Collard. 1998. Entre Atlantique et Méditerranée : grand tétras, lagopède, perdrix grise des Pyrénées et marmotte. Statut territorial et évolution. *Bulletin mensuel de l'Office National de la Chasse*. 231 : 16-23.
- Ménoni, E. & O. Duriez. (2008). Le Grand tétras *Tetrao urogallus* dans les Pyrénées : historique et statut actuel. *Ornithos* 15(4) : 272-281.
- Ménoni, E., C. Novoa, C. Berducou, J. Canut, M. Mossoll-Torres, M. Monta, S. Marin, J. Piqué, C. D., & J. A. Gil Gallus. (2004). Évaluation transfrontalière de la population de Grand tétras des Pyrénées. *Faune sauvage* 263 : 19-24.
- Mollet, P., M. Kéry, B. Gardner, G. Pasinelli, and J. A. Royle. 2015. Estimating Population Size for Capercaillie (*Tetrao urogallus* L.) with Spatial Capture-Recapture Models Based on Genotypes from One Field Sample. *Plos one* 0(6).
- Montadert, M. 2012. Partial recovery of capercaillie in French Jura Mountains. *Grouse news* 44 : 9-15.
- Montadert, M. 2013. Statut actuel et variation d'abondance du Grand tétras (*Tetrao urogallus*) dans le massif jurassien français. *Alauda* 81 : 1-18.
- Moreira Boman, C., A. Pettersson. 2020. Limiting factors in capercaillie (*Tetrao urogallus*) conservation. *Kandidatarbeten in skogsvetenskap. Fakulteten for skogsvetenskap*.
- Moss, R. 1994. Decline of capercaillie (*Tetrao urogallus*) in Scotland. *Gibier Faune Sauvage, Game and Wildlife*. 11:(217-222).
- Mossoll-Torres, M., and E. Ménoni. 2006. The Capercaillie in Andorra: a population in a good health! *Grouse news* 32 : 18-23.
- O.G.M. 2011. Évolution de la répartition communale du petit gibier de montagne en France, décennie 2000-2009. *Faune sauvage* 290-Supplément : 16 pp.

Obeso, J. R., and M. J. Bañuelos. 2003. El urogallo (*Tetrao urogallus cantabricus*) en la Cordillera Cantabrica. Ministerio de Medio Ambiente. Naturaleza y Parques Nacionales. Edition. Oviedo.

Office National de la Chasse. 1977. Enquête nationale sur la situation du grand tétras. Bulletin Mensuel de l'ONC. Supplément 4 : 23 pp.

Peterhofs, E., G. Bagrade, G. Done, M. Lukins, A. Ornicans, J. Ozolins, D. Pilate, D.-E. Rungis, J. Suba, and A. Zunna. 2019. Suitability of capercaillies (*Tetrao urogallus*) as an umbrella species for Latvian boreal forests. *in* Proceedings of 34th IUGB Congress 2019.

Rassati, G. 2009. Il Gallo cedrone (*Tetrao urogallus*) in due area campione delle Alpi Carniche (Friuli-Venezia Giulia). *Riv. ital. Orn.*, Milano 79:49-57.

Rimbeuf, M. 2020. Réactualisation de l'aire de répartition de Grand Tétras dans le massif vosgien. Stage de Master 2, Groupe Tétras Vosges. 58 p.

Robles, L., F. Ballesteros, and J. Canut. 2007. El urogallo en España, Andorra y Pirineos franceses. Situation actual (2005). SEO/Birdlife, Madrid.

Schroeder, M. A., R. K. Baydack, S. A. harmon, C. A. Hagen, D. M. Davis, S. K. Sherrod, S. deMaso, R. W. Hoffman, T. Z. Riley, J. B. Haufler, and R. R. Manes. 2004. The North American Grouse Management Plan: a prospectus. North American Grouse partnership.

Storch, I. 2000. Grouse status survey and conservation action plan 2000-2004. U.K. and the world pheasant association, Switzerland and Cambridge.

Storch, I. 2006. An overview to the conservation status of grouse in Europe. Pages 15-22 *in* Proceedings of Alpines Grouses Conservation and Management experiences across Europe : 15-22.

Storch, I. 2010. Grouse : Status survey and conservation action plan 2006-2010. IUCN Species Survival Commission.

Vázquez, J. F., P. T. J. Albornoz, and A. Domínguez. 2013. Census and effective population size of the endangered Cantabrian capercaillie (*Tetrao urogallus*) estimated from non-invasive samples. *Grouse news* 46 : 12-26.

Viht, E. 1994. Metsislaste seirest eestis (in estonian, with english summary) : On the monitoring of the tetraonids in Estonia. *Hirundo* 2 : 59-65.

Viht, E., and T. Randla. 2002. Metsis eesti asurkonna seisund ja säilimise väljavaated. The capercaillie: status of the estonian population and its survival prospects. (In estonian and english). *Hirundo Supplementum* 5 : 50 p.

Watson, A., and R. Moss. 2004. Grouse : The natural history of British and Irish species. HarperCollins, London, UK. 529pp.

Zawadzka, D., J. Zawadzki, and M. Piotrowska. 2007. 1st International Conference, "Conservation of the Forest Grouse" in Poland. *Grouse news* 34 : p.18.

Biologie-écologie

Ahlen, P.-A., T. Willebrand, S. K., and M. Hörnell-Willebrand. 2013. Survival of female Capercaillie *Tetrao urogallus* in northern Sweden. *Wildlife Biology* : 368-373.

Bortchevski et al. 2003. Does fragmentation by logging reduce grouse reproductive success in boreal forests ? *Wildlife Biology*, 9, 275-283.

Eliassen, S., and P. Wegge. 2007. Ranging behaviour of male capercaillie *Tetrao urogallus* outside the lekking ground in spring. *Journal of Avian Biology* 38 : 37-43.

Groupe Tétras Vosges. Synthèses annuelles des suivis de la population de Grand Tétras par pièges photographiques dans les Vosges, 2018 à 2022.

Hjorth, I. 1970. Reproductive behaviour in Tetraonidae with special reference to males. *Viltrevy* 7 : 183-596.

Holmstad, P. R., P. J. Hudson, V. Vandvik, and A. Skorping. 2005. Can parasites synchronise the population fluctuations of sympatric tetraonids ? Examining some minimum conditions. *Oikos* 109 : 429-434.

HUDSON, P. J., and D. NEWBORN. 1990. Brood defence in a precocial species: variations in the distraction displays of red grouse, *Lagopus lagopus scoticus*. *Animal Behaviour* 40 : 254-261.

Lebreton, P., and I. Girard. 2006. Etude écologique des galliformes de montagne : compléments méthodologiques. *Travaux Scientifiques du Parc National de la Vanoise* XXIII : 147-158.

Leclercq, B., and E. Ménoni. 2018a. Le Grand tétras. *Biotope*, Mèze, 352 pp.

Liukkonen, T. 2007. Observations on the flocking and mobility of Capercaillie (*Tetrao urogallus*)- hunters'fairytales or true observations? *Ethology Ecology & Evolution* 19 : 245-255.

Liukkonen, T., J. Bisi, and S. Kurki. 2006. Observations on capercaillie *Tetrao urogallus* flocking and mobility- facts or foiction? *Grouse news* 32 : 7-11.

Ménoni, E. 1990. Caquètements et territorialité des poules de grand tétras au printemps dans les Pyrénées. (in french, with english summary: spring cackle calls and territoriality in hens capercaillie in the Pyrénées). *Acta Biologica Montana* 10:(63-82).

Ménoni, E., M. Clemente, G. Chasseraud, & L. C. E. C. Berducou. (2011). Enseignements écologiques et éthologiques tirés du comportement anormal d'un Grand tétras *Tetrao urogallus* en vallée d'aspe (pyrénéesatlantiques). *Alauda* (3) 79 : 199-206.

Spidso, K., P. Wegge, and T. Storaas. 1984. Renesting in Capercaillie in southern Norway. Pages 278-290 *in* Proceedings of 4rd International Symposium on Grouse. Lovel T.W.I. World Pheasant Association ed.:278-290.

Storaas, T., P. Wegge, and L. Kastdalen. 2000. Weight-related renesting in capercaillie *Tetrao urogallus*. *Wildlife Biology* 6 : 299-303.

Wegge, P. 1985. Spacing pattern and habitat use of capercaillie hens in spring. Pages 261-277 *in* Proceedings of 3th International Symposium on Grouse. 261-277.

Wegge, P., S. Eliassen, M. H. Fine, and M. Odden. 2005a. Social interactions among Capercaillie *Tetrao urogallus* males outside the lek during spring. *Ornis Fennica* 82 : 147-154.

Wegge, P. and Rolstad J., 2011, Clearcutting and eurasian boreal grouse : Long term monitoring reveals unexpected effects on their population performances. Forest Management, 261.

Fonctionnement et dynamique des populations

Abrahams, C. 2019. Comparison between lek counts and bioacoustic recording for monitoring Western Capercaillie (*Tetrao urogallus* L.). Journal of Ornithology. <https://doi.org/10.1007/s10336-019-01649-8>:13pp.

Aleix-Mata, G., B. Adrados, M. Boos, E. Marty, P. Mourieres, G. Tucat, N. Thion, M. Mossoll-Torres, J. M. Pérez, and A. Sánchez. 2020. Comparing methods for estimating the abundance of Western Capercaillie *Tetrao urogallus* males in Pyrenean leks: singing counts versus genetic analysis of non-invasive samples. Bird Study:6 pp.

Augustine, B. C., M. Kéry, J. Olano-Marin, P. Mollet, G. Pasinelli, and C. Sutherland. 2019. Sex-specific population dynamics and demography of capercaillie (*Tetrao urogallus* L.) in a patchy environment. bioRxiv.

Baguette, M. (2004). The classical metapopulation theory and the real, natural world : a critical appraisal. Basic and Applied Ecology 5 : 213-224.

Bal, G., L. Bacon, E. Ménoni, C. Calenge, A. Millon, and A. Besnard. 2021. Modélisation de la dynamique du grand tétras des Pyrénées françaises pour sa gestion adaptative. Technical report MNHN et OFB.

Belovski, G. E., J. A. Bissonette, R. D. Dueser, T. C. Edwards, C. M. Luecke, M. E. Ritchie, J. B. Slade, and F. H. Wagner. (1994). Management of small populations : concepts affecting the recovery of endangered species. Wildlife society bulletin 22 : 307-316.

Borchtchevski, V. 1993. Population biology of the capercaillie. Principles of the structural organisation. 268 pp. in Russian with english summary and figures. Moscou edition. Moscou.

Borchtcheski, V. and M. Moss. 2013. Age structure of Capercaillie males (*Tetrao urogallus*) in NW Russia may reflect two-way movements. A hypothesis. Ornis fennica 90 : 15 pp.

Cayuela, H., J. G. Prunier, M. Laporte, J. Gippet, L. Boualit, F. Preiss, A. Laurent, F. Foletti, and G. Jacob. 2019b. Demography, genetic, and extinction process in a spatially structured population of lekking bird. in press.

Gée, A., M. Sarasa, and O. Pays. 2018. Long-term variation of demographic parameters in four small game species in Europe: opportunities and limits to test for a global pattern. Animal Biodiversity and Conservation 41.

Grimm, V., and I. storch. 2000. Minimum viable population size of capercaillie *Tetrao urogallus*: results from a stochastic model. Wildlife Biology 6:219-225.

Guixé, D., and X. Florensa. 2020. Protocolo de seguimiento del urogallo común (*Tetrao urogallus*) con métodos acústicos y de fototrampeo. Centro de Ciencia y Tecnología Forestal de Cataluña.

Isomursu, M., O. Rätti, P. Helle, and T. Hollmen. 2006. Sex and age influence intestinal parasite burden in three boreal grouse species. Journal of Avian Biology 37:516-522.

Jaako, A., A. Putala, and A. Marjakangas. 2005. Factors affecting nesting success in black grouse. (in finnish, with english summary. Teeren pesän säilymiseen vaikuttavista tekijöistä). Suomen riista 51:44-50.

Leclercq, B. 1988. Ecologie et dynamique des populations du grand tétras (*Tetrao urogallus major L.*) dans le Jura français. PhD., Université de Bourgogne, Dijon.

Ludwig, G. X., R. V. Alatalo, P. Helle, H. Linden, J. Lindstrom, and S. H. 2006. Short- and long-term population dynamical consequences of asymmetric climate change in black grouse. Proceeding of the Royal Society Bulletin 273:2009-2016.

Ménoni, E. 1991. Ecologie et dynamique des populations du grand tétras dans les Pyrénées avec des références spéciales à la biologie de la reproduction chez les poules. Quelques applications à sa conservation. PHD, Université Paul Sabatier., Toulouse.

Ménoni, E. & C. Novoa. 1988. Apport de l'analyse des tableaux de chasse dans la connaissance des populations de Grand tétras (*Tetrao urogallus L.*). Gibier Faune Sauvage 5(3) : 255-272.

Milhau, B. et al. 2023. Evaluation de la stratégie nationale en faveur du Grand Tétras (2012-2022). OGM.

Moss, R., and J. Oswald. 1985. Population dynamics of Capercaillie in a North-east Scottish glen. *Ornis Scandinavica* 16:229-238.

Moss, R., and D. N. Weir. 1987. Demography of capercaillie *Tetrao urogallus* in North-East Scotland. III. Production and recruitment of young. *Ornis scandianavica* 18 : 141-145.

Moss, R., and A. Watson. 1990. Breeding success and movement as determinants of breeding density in tetraonid birds. Pages 85-92 in Proceedings of XIX IUGB Congress.85-92.

Moss, R., N. Picozzi, R. W. Summers, and D. Baines. 2000. Capercaillie *Tetrao urogallus* in Scotland-demography of a declining population. *Ibis* 142:(259-267).

Moss, R., J. Oswald, and D. Baines. 2001. Climate change and breeding success : decline of the capercaillie in Scotland. *Journal of animal ecology*. 70 : 47-61.

Moss, R., N. Picozzi, and D. C. Catt. 2006. Natal dispersal of capercaillie (*Tetrao urogallus*) in northeast Scotland. *Wildlife Biology* 12:227-232.

PALANQUE, D. 1999. Evolution des effectifs de grands tétras (*Tetrao urogallus*) dans le massif vosgien. *Gibier Faune Sauvage, Game and Wildlife*. 16 : 225-249.

Rajala, P., and H. Linden. 1974. Finnish tetraonid populations in August 1972 compared with the period 1963/1971 according to route census. *Suomen Riista* 25 : 56-63.

Ranta, E., J. Lindstrom, H. Linden, and P. Helle. 2008. How reliable are harvesting data for analyses of spatio-temporal population dynamics? *Oikos* 117 : 1461-1468.

Rotelli, L., and M. Bocca. 2004. Proposte per la gestione dei galliformi sulle Alpi occidentali italiane. Pages (63-76) in Proceedings of Atti del Seminario "Esperienze di gestione dei Galliformi di Montagna con particolare riferimento alla programmazione venatoria". 63-76.

Sachot, S. 2002. Viability and management of an endangered capercaillie (*Tetrao urogallus*) metapopulation. Ph. Thesis, Université de Lausanne., Lausanne.

Sachot, S., B. Leclercq, and M. Montadert. 2002. Population trends of capercaillie (*Tetrao urogallus*) in the Jura mountains between 1991 and 1999. Game and Wildlife Science. 19 : 41-54.

Sachot, S., N. Perrin, and N. Cornelis. 2006. Viability and management of an endangered Capercaillie (*Tetrao urogallus*) metapopulation in the Jura Mountains, western Switzerland. Biodiversity and Conservation 15:2017-2032.

Saniga, M. 1999. Population dynamics of Capercaillie *Tetrao urogallus* on leks in Central Slovakia in the period 1981-1997. VogelWelt 120:235-240.

Saniga, M. 2012. Population dynamics of Capercaillie *Tetrao urogallus* on leks in Central Slovakia in the period 1981-2012. Grouse news 44:5-9.

Selas, V. 2000. Population dynamics of Capercaillie in relation to bilberry production in southern Norway. Wildlife Biology, 6, 1-11.

Selas, V., et al. 2021. Bilberry seed production explains spatio temporal syncronicity in bank vole population fluctuations. Ecological Research, 36, 409-419.

Souchay, G., A. Besnard, C. Perrot, C. Jakob, and F. Ponce. 2018. Anthropic and natural factors drive variation of survival in a fast galliform species in southern France. Wildl. Biol. in press : 39 pp.

Schwartz, T., A. Besnard. 2022. Viabilité de la population de Grand Tétras du Jura. Etude de différents scénarios de renforcement. Plan National d'Action en faveur du Grand Tétras, déclinaison Jura. CEFE. A. Rocha.

Schwartz, T., A. Besnard. 2021. Modélisation de la viabilité de la population de Grand Tétras (*Tetrao urogallus*) du massif des Vosges selon différents scénarios de renforcement. Etude de faisabilité. Plan National d'Action en faveur du Grand Tétras, déclinaison Vosges. CEFE. A. Rocha. PNRBV. GTV.

Wegge, P. 1980. Distorted sex ratio among small broods in a declining Capercaillie population. Ornis Scandinavica 11:106-109.

Wegge, P., and E. Rolstad. 2012. GPS satellite telemetry: a promising tool for grouse research. Grouse news 43:9-11.

Wegge, P., R. Moss, J. Rolstad. 2022. Annual variation in breeding success in boreal forest grouse : Four decades of monitoring reveals bottom-up drivers to be more important than predation. Ecology and Evolution. DOI : 10.1002/ece3.9327.

Prédation

Alert, B. and I. Storch. (2009). Does predation affect lekking behaviour in grouse ? A summary of field observations. Grouse news 38 : 30-42.

Andren, H. (1992). Corvid density and nest predation in relation to forest fragmentation : a landscape perspective. Ecology 73(3): 794-804.

Andren, H., P. Angelstam, E. Lindström, and P. Widen. (1985). Differences in predation pressure in relation to habitat fragmentation : an experiment. Oikos 45 : 273-277.

- Angelstam, P. (1986). Predation on ground nesting birds' nests in relation to predator densities and habitat edge. *Oikos* 47(3): 365-373.
- Angelstam, P., E. Lindström, and P. WIDEN. (1985). Synchronous short-term population fluctuations of some birds and mammals in Fennoscandia – occurrence distribution. *Holarctic Ecology* 8 : 285-298.
- Angelstam, P., E. Lindström, and P. Widen. (1984). Role of predation in short-term population fluctuations of some birds and mammals in Fennoscandia. *Oecologia* 62 : 199-208.
- Anker Ims, R., J. Rolstad, and P. Wegge. (1993). Predicting space use responses to habitat fragmentation : can voles *Microtus oeconomus* serve as an experimental model system (EMS) for Capercaillie grouse *Tetrao urogallus* in boreal forest ? *Biological Conservation* 63 : 261-268.
- Armstrong, E. A. (1955). Distraction display and the human predator. *Ibis* 98 : 641- 654.
- Baines, D., R. Moss, and D. Dugan. (2004). Capercaillie breeding success in relation to forest habitat and predator abundance. *Journal of Applied Ecology* 41 : 59-71.
- Baines, D., R. B. Sage, and M. M. Baines. (1994). The implication of red deer grazing to ground vegetation and invertebrate communities of Scottish native pinewoods. *Journal of applied ecology*. 31 : 776-783
- Beasley, J. C., Z. H. Olson, W. S. Beatty, G. Dharmarajan, and O. J. Rhodes. (2013). Effects of culling on mesopredator population dynamics. *Plos One* 6 (11).
- Beasom, S. L. (1974). Intensive short-term predator removal as a game management tool. Thirty-ninth north American wildlife and natural resources conference., Denver, Wildlife management institute.
- Cas, M. 2008. Disturbance at endangered capercaillie leks in Slovenia caused by predation and other reasons in the period between 1984-2000. *Acrocephalus* 29 : 6pp.
- Drillon, V. 1995. Bilan du piégeage expérimental mené pendant cinq ans sur le massif pilote de la Haute-Meurthe. Pages 137-150 in *Proceedings of Le Grand tétras, question de survie* : 137-150.
- Fernández-Olalla, M., A. Martínez-Abraín, J. Canut, D. García-Ferré, I. Afonso, and L. Mariano-González. 2012. Assessing different management scenarios to reverse the declining trend of a relict capercaillie population: A modelling approach within an adaptive management framework. *Biological Conservation* 148:79-87.
- Fletcher, K., A. N. Hoodless, and D. Baines. 2013. Impacts of predator abundance on red grouse *Lagopus lagopus scotica* during a period of experimental predator control. *Wildlife Biology* 19: 248-256.
- Giraudoux P., Leclercq B. & Desbrosses R. (1997). Cycles démographiques du Campagnol terrestre (*Arvicola terrestris*) et reproduction et cinétique des populations de tétraonidés du massif du Risoux (Jura). *Gibier Faune Sauvage, Game & Wildlife* 14: 31- 47.
- HELDIN, J. O. 2000. Population trends and harvest management of pine marten *Martes martes* in Scandinavia. *Wildl.Biol.* 6.
- HUDSON, P. J., A. P. DOBSON, and D. NEWBORN. 1992. Do parasites make prey vulnerable to predation? Red grouse and parasites. *Journal of Animal.Ecology* 61:681-692.
- Jahren, T. et al. 2017. Declining reproductive output in Capercaillie and black grouse : 16 countries and 80 years. *Animal Biology*, 363-400.

Jahren, T. 2017. The role of nest predation and nest predators in population declines of capercaillie and black grouse. Inland Norway University of Applied Sciences, Evenstad.

Kammerm   J.-L., I. Storch. 2019. "Predation, predator control and grouse populations : a review. Wildlife Biology, wlb 00464.

K  mmerle, i.-L., J. Coppes, R. Suchant, and I. Storch. 2020. From research to practice : Effectiveness of restricted-area fox culls for capercaillie conservation (in german, with english summary and figures). Natur und Landshaft 12 : 525-531.

Kauhala, K., P. Helle, and E. Helle. 2000. Predator control and the density and reproductive success of grouse populations in Finland. ECOGRAPHY 23 : 161-168.

LECLERCQ, B., R. DESBROSSES, and P. GIRAUDOUX. 1997. Cycles d  mographiques du campagnol terrestre (*Arvicola terrestris*), et reproduction et cin  tique des populations de *tetraonidae* du massif du Risoux (Jura). Gibier Faune Sauvage, Game Wildlife 14 : 31-47.

LINDEN, H. 1988. Latitudinal gradients in predator-prey interactions, cyclicity and synchronism in voles and small game populations in Finland. Oikos 52:341-349.

Lindstr  m, E. R., n. H. Andre, P. Angelstam, G. Cederlund, B. H  rnfeldt, L. J  derberg, P.-A. Lemnell, B. Martinsson, K. Sk  ld, and J. E. Swenson. 1994. Disease reveals the predator: sarcoptic mange, red fox predation, and prey populations. Ecology 75 : 1042-1049.

Ludwig, G. X., R. V. Alatalo, P. Helle, I. Koljola, and H. Siitari. 2008. Large carnivore reinvasion and the future of Finnish Forest Grouse. *in* Proceedings of XI International Symposium on Grouse.

Marcstr  m, V., R. E. Kenward, and E. Engren. 1988. The impact of predation on boreal tetraonids during vole cycles : an experimental study. Journal of Animal Ecology 57 : 859-872.

M  noni, E., and J. Bougerol. 1993. Capercaillie population in forest fragmented by topography and human activities in the french Pyrenees. Pages (148-159) *in* Proceedings of XXI congress I.U.G.B. 2 : 148-159.

M  NKK  NEN, M., M. HUSBY, R. TORNBERG, P. HELLE, and R. L. THOMSON. 2007. Predation as a landscape effect: the trading off by prey species between predation risks and protection benefits. *Journal of Animal Ecology* 76 : 619-629.

M  NKK  NEN, M., TORNBERG. R, and P. VAEISAENEN. 2000. Goshawks may reduce predation rates on birds'nests. Suomen riista 46 : 27-36.

Moreno-Opo, R., I. Afonso, J. Jim  nez, M. Fern  ndez-Olalla, J. Canut, D. Garc  a-Ferr  , J. Piqu  , F. Garc  a, J. Roig, J. Mu  oz-Igualada, L. Mariano-Gonz  lez, and J. V. L  pez-Bao. 2015. Is It Necessary Managing Carnivores to Reverse the Decline of Endangered Prey Species? Insights from a Removal Experiment of Mesocarnivores to Benefit Demographic Parameters of the Pyrenean Capercaillie. PLoS ONE DOI : 10.1371/journal.pone.0139837.

Moreno-Opo, R. et al. Ann  e ? Insights from a removal experiemnt of mesocarnivores to benefit demographic parameters of Pyrenean Capercaillie. PLoS ONE DOI 10.1371/journal.pone.0139837.

Oja, R., E. Soe, H. Valdmann, and U. Saarma. 2017. Non-invasive genetics outperforms morphological methods in faecal dietary analysis, revealing wild boar as a considerable conservation concern for ground-nesting birds. PLoS ONE 12.

Pasanen-Mortensen1, M., M. Pyykönen, and B. Elmhagen. 2013. Where lynx prevail, foxes will fail – limitation of a mesopredator in Eurasia. *Global Ecology and Biogeography* 22:868-877.

Redpath, S. M., B. E. Arroyo, E. M. Leckie, P. Bacon, N. Bayfield, R. J. Gutierrez, and S. J. Thirgood. 2004. Using Decision Modeling with Stakeholders to Reduce Human-Wildlife Conflict: a Raptor-Grouse Case Study. *Conservation Biology* 18:350-359.

Roda, F., and Roda.J-M. 2016. Foraging traces as an indicator to monitor wild boar impacton ground nesting birds. <hal-01306234>.

RUETTE, S., N. LIEURY, M. ALBARET, and J.-P. Arnauduc. 2015. Evolution des populations de renard en France, Analyse des suivis réalisés par comptages nocturnes (2004-2013). *Faune sauvage* 306:37-42.

Saniga, M. 2002. Nest loss and chick mortality in capercaillie (*Tetrao urogallus*) and hazel grouse (*Banasa bonasia*) in West Carpathians. *Folia Zoologica* 51 : 205-214.

Signorell, N., S. Wirthner, P. Patthey, R. Schranz, L. Rotelli, and R. Arlettaz. 2010. Concealment from predators drives foraging habitat selection in brood-rearing Alpine black grouse *Tetrao tetrix* hens: habitat management implications. *Wildl.Biol.* 16.

Smedshaug, C. A. 2001. Was high grouse bag in early 20th century Norway due to a program for extermination of small game predators? *Ecography* 24 : 579-587.

Stahl, P., and P. Migot. 1993. L'impact des prédateurs sur le petit gibier. Une revue des enlèvements expérimentaux de prédateurs. Pages 21-35 in Proceedings of Colloque préddation et gestion des prédateurs. 21-35.

Storaas, T., and P. Wegge. 1984. High nest losses in Capercaille and Black grouse in Norway. Lovel T.W.I. World Pheasant Association ed. Pages 278-290 in Proceedings of 3rd International Symposium on Grouse.278-290.

Storaas, T. 1987. A comparaison of losses in artificial and naturally occuring capercaillie nests. *Journal of wildlife management* 52:123-126.

STORAAS, T., L. KASTDALEN, and P. WEGGE. 1999. Detection of forest grouse by mammalian predators. A possible explanation for high brood losses in fragmented landscapes. *Wildlife Biology* 5:187-192.

Storaas, T. 2001. Forest fragmentation increases brood mortality of grouse by mammalian predators: a hypothesis. *Suomen riista* 47:86-93.

Storch, I. 2018. Is the decline of grouse in central Europe related to landscapes favoring predators? in Proceedings of XIVth International Symposium on Grouse.

Storch, I., E. Woitke, and S. Krieger. 2005. Landscape-scale edge effect in predation risk in forest-farmland mosaics. *Landscape Ecology* 20 : 927-940.

Sulkava, S., K. Huhtala, P. Rajala, and R. Tornberg. 2003. Prey selection by the golden eagle between tetraonid species and their sexes in Finland during 1957-1997. in finnish with english summary: Kotlan saaliinvalinta metsäkanalintujen ja niiden sukupuolten kesken Suomessa 1957-97. *Suomen riista* 49:(44-54).

Summers, R. W., J. Willi, & J. Selvidge. 2009. Capercaille *Tetrao urogallus* nest loss and attendance at Abernethy Forest, Scotland. *WildlifeBiology* 15: 319-327

Summers, R. W., D. Dugan, and R. Proctor. 2010. Numbers and breeding success of Capercaillies *Tetrao urogallus* and Black Grouse *T. tetrix* at Abernethy Forest, Scotland. *Bird Study*:437-446.

Summers, R. W., R. E. Green, R. Proctor, D. Dugan, D. Lambie, R. Moncrieff, R. Moss, and D. Baines. 2004. An experimental study of the effects of predation on the breeding productivity of capercaillie and black grouse. *Journal of Applied Ecology* 41:(513-525).

Thiel, D., E. Ménoni, J. F. Brenot, and L. Jenni. 2007. Effects of recreation and hunting on flushing distance of capercaillie. *Journal of Wildlife Management*, 71:1784-1792.

Thiel, D. K., and K. Bollmann. unpublished. Effects of forest size and forest fragmentation on predation rate of artificial ground nest in the Swiss Alps. 1-20.

Tornberg, R., E. Korpimäki, S. Jungell, and V. Reif. 2005. Delayed numerical response of goshawks to population fluctuations of forest grouse. *Oikos* 111:408-415.

Tornberg, R., A. Linden, P. Byholm, E. Ranta, J. Valkama, P. Helle, and H. Linden. 2013. Coupling in goshawk and grouse population dynamics in Finland. *Oecologia* 171:863-872.

Tornberg, R., V. Reif, and E. Korpimäki. 2012. What Explains Forest Grouse Mortality: Predation Impacts of Raptors, Vole Abundance, or Weather Conditions? *International Journal of Ecology Article ID 375260*, doi:10. 1155/2012/375260 : 10 pages.

TORNBERG, R. 2000. EFFECT OF CHANGING LANSCAPE STRUCTURE ON THE PREDATOR-PREY INTERACTION BETWEEN GOSHAWK AND GROUSE. PhD, Oulu, Oulu.

TORNBERG, R. 2005. The response of the goshawk *Accipiter gentilis* to changing grouse *tetraonidae* sp. populations. Pages 160-167 *in* Proceedings of Workshop of Kostomuksha.160-167.

Valkama, J., E. Korpimäki, A. B., P. Beja, V. Bretagnolle, E. Bro, R. Kenward, S. Manosa, R. S.M., S. Thirgood, and V. J. 2005. Birds of prey as limiting factors of gamebird populations in Europe: a review. *Biological review*. 80:(171-203).

Wegge, P., H. Ingul, V. O. Pollen, E. Halvorsrud, A. V. Sivkov, and H. O. 2012. Comparing predation on forest grouse nests by avian and mammalian predators in two contrasting boreal forest landscapes by the use of artificial nests. *Ornis Fennica* 89:XXX-XXX (online early).

Wegge, P., B. B. Larsen, I. Gjerde, L. Kastdalen, J. Rolstad, and T. Storaas. 1987. Natural mortality and predation of adult capercaillie in southeast Norway. Pages (50-56) *in* Proceedings of 4th int. Grouse Symposium. 50-56.

Willebrand, T., S. Willebrand, T. Jahren, and V. Marcström. 2017. Snow tracking reveals different foraging patterns of red foxes and pine martens. *Mammalian research*.

Zubic, G. 2009. Capercaillie *Tetrao urogallus* in Serbia - principal threats and conservation measures. *Grouse news* : 32-33.

Zubic, G. 2010. Density and vulnerability of capercaillie *Tetrao urogallus* in the area of Vitoroga in the Republika Srpska. *Grouse news* : p. 10-11.

Habitat

Alvarez, E., P. Alvarez, B. F., M. J. Banuelos, E. Blanco, M. Calvo, J. Del Campo, J. Camprodon, F. J. Ezquerra, I. C. Fernandez, J. Garrote, L. M. Gonzalez, F. Gonzalez, F. J. Jimenez, M. Lucas de, A. J. Lucio, J. Majada, V. Mateos, J. I. Molina, J. R. Obeso, R. Otero, C. J. Pollo, F. J. Purroy, J. Reque, L. Robles, A. Rossignoli, and F. Sevilla. 2005. Manual de Conservacion y Manejo del Habitat del Urogallo Cantabrico. Direccion General para la Biodiversidad. Ministerio de Medio Ambiente. Madrid.

Andren, H. 1994. Effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat : a review. *Oikos* 71(3): 355-366.

Andren, H. 1996. Population responses to habitat fragmentation : statistical power and the random sample hypothesis." *Oikos* 76(2): 235-242.

Angelstam, P. 2004. Habitat thresholds and effects of forest landscape change on the distribution and abundance of Black grouse and Capercaillie. *Ecological bulletins* 51 : 173-187.

Angelstam, P., V. M. Anufriev, L. Balsiauskas, A. K. Blagovidov, S.-O. Borgegard, S. J. Hodge, P. Majewski, S. V. Ponomarenko, E. A. Shvarts, A. A. Tishkov, L. Tomialojc, and T. Wesolowski. 1997. Biodiversity and sustainable forestry in European forests : how east and west can learn from each other. *Wildlife society bulletin* 25 : 38-48.

Banuelos, M. J., M. Quevedo, and J. R. Obeso. 2008. Habitat partitioning in endangered cantabrian Capercaillie *Tetrao urogallus cantabricus*. *Journal of Ornithology* 149 : 245-252.

Barikmo, J., P. Osbak, and O.-K. Bokseth. 1984. Distances of movements and spacing behaviour for Capercaillie broods (*Tetrao urogallus*). In proceedings of third international grouse symposium : 291-300.

Berchtold, M., L. Rotelli, and I. Storch. 2010. Habitat selection and spacing behavior of Capercaillie in the Italian Alps : a project in progress. *Grouse news* 40 : 20-21.

Berducou, C. (1996). Gestion forestière et Grand tétras-Pyrénées. ONF édition.

Bérion E., Leclercq B. Et Montadert M. 2005. Influence de la structure et de la composition de la végétation sur l'abondance du Grand tétras *Tetrao urogallus* L. dans le Massif jurassien français. In « Les capacités d'accueil des forêts pour le Grand tétras ». Doc. interne Groupe Tétras Jura, contrat DIREN Franche Comté : 9-17.

Bernard, B. (2006). La gestion locale des forêts à l'épreuve du tétras. *Terrain et Travaux* 11 : 16 pp.

Betton, T., C. Novoa, E. Ménoni, K. Foulché, and F. Sèbe. (2017). Acoustic Survey for Biodiversity Response Measurement to Habitat Restoration. *Conservation Biology* : 13 pp.

Betton, T., K. Foulché, E. Ménoni, C. Novoa, F. Nicolè, N. Mathevon, and F. Sèbe. 2018. Acoustic monitoring of biodiversity response to habitat restoration. *in Proceedings of XIVth International Symposium on Grouse*.

Blanco-Fontao, B., A. Fernandez-Gil, J. R. Obeso, and M. Quevedo. (2010). Diet and habitat selection in cantabrian Capercaillie (*Tetrao urogallus cantabricus*) : ecological differentiation of a rearedge population. *Journal of Ornithology* 151 : 269-277.

Blanco-Fontao, B., J. R. Obeso, M. J. Banuelos, and M. Quevedo. (2011). Habitat partitioning and molting site fidelity in *Tetrao urogallus cantabricus* revealed through stable isotopes analysis. *Journal of Ornithology*.

Bollmann, K., P. Weibel, and R. Graf. (2005). An analysis of central Alpine Capercaillie spring habitat at the forest stand scale. *Forest Ecology and Management*. 215 : 307-318.

Borchtchevski, V. and R. Moss. 2008. Factors affecting the spring distribution of Capercaillie (*Tetrao urogallus*) and Black grouse (*T. tetrix*) in habitats of largely-intact taiga in NW Russia. *News of Game Management* (Moscow). (In Russian with English summary) 5(2): 138-161.

Borchcheski, V. (in prep.). Spring habitat use by sympatric populations of Capercaillie, Tetrao urogallus, and Black grouse, *T. Tetrix*, in native taiga landscape of N-W Russia : 1-20.

Braunisch, V. 2006. A habitat and habitat-connectivity model for Capercaillie *Tetrao urogallus* in the Black Forest as a tool for sustainable conservation planning (a PhD project in progress). *Grouse news* 31 : 9-11.

Braunisch, V., K. Bollmann, G. R.F., and A. H. Hirzel. 2008. Living on the edge modelling habitat suitability for species at the edge of their fundamental niche. *Ecological modelling* 214 : 153-167.

Braunisch, V. and R. Suchant. 2007. A model for evaluating the “habitat potential” of a landscape for Capercaillie *Tetrao urogallus* : a tool for conservation planning. *Wildlife Biology* 13 (suppl.) : 21-23.

Braunisch, V., and R. Suchant. 2008. Using ecological forest site mapping for long-term habitat suitability assessments in wildlife conservation—Demonstrated for capercaillie (*Tetrao urogallus*). *Forest Ecology and Management* 256:1209-1221.

Campión, D., and J. Camprodón. 2011. Silvicultura. Estrato arbóreo. Pages 52-72 in J. Canut, García-Ferré, D. & Afonso, I. Dirección General del Medio Natural y Política Forestal. Ministerio de Medio Ambiente, Medio Rural y Marino. Serie manuales de gestión de especies amenazadas., editor. *Manual de conservación y manejo del urogallo pirenaico*. Madrid.

Camprodón, J., D. G. Coromines, D. Garcia-Ferré, and P. Omedes-Perez. 2016. Como lograr un mejor habitat para el urogallo en los pirineos. *Quercus* 366 : 14-20.

Cantegrel, R. 2013. Evaluation patrimoniale des populations autochtones de pin à crochets (*Pinus uncinata* Ramond) aux Pyrénées. *Les Dossiers Forestiers-ONF* 25 : 159 pp.

Canut, J. 2001. Gallinaceas de montaña (perdiz pardilla, lagopodo alpino y urogallo) y gestión forestal. Pages 291-306 in J. Camprodón, and E. Plana-Bach, editors. *Conservation de la biodiversidad y gestión forestal. Su aplicación en la fauna vertebrada*. barcelona.

Canut, J., D. García-Ferré, and I. Afonso. 2011. *Manual de conservación y manejo del hábitat del urogallo pirenaico*. Madrid.

Čas, M. 2007. Fluctuations in the capercaillie *Tetrao urogallus* (L.) population in relation to past land use and forest structures in the south-east Alps. *Grouse news* 33 : 20.

Charbonnier, G., T. Chevalier, F. Preiss. 2013. Diachronie des milieux ouverts sur le massif de Ventron (88). Groupe Tétrias Vosges. 11p.

DESCHAMPS, J., F. ROUTIER, E. MENONI, and L. LARRIEU. 2017. Capercaillie: is it a meaningful indicator for French central Pyrenees forest biodiversity? in Proceedings of 33e Congrès IUGB.- Montpellier, France.

Devau, B., and M. Catusse. 1988. Habitats utilisés par le Grand Tétras (*Tetrao urogallus aquitanicus*) dans la forêt pyrénéenne française en hiver et au printemps. Pages 69-84 in Proceedings of Colloque Galliformes de Montagne.69-84.

Fattebert, P., S. Sachot, N. Perrin, and C. Neet. 2003. Model development for capercaillie (*Tetrao urogallus*) habitat in the Jura mountains (western Switzerland). *Game and Wildlife Science* 20:(195-210).

Gabrielli, J., E. Ménoni, B. Milhau, M. Le Vaillant, S. Pascal, A. Berceau, and K. Foulché. 2015. Prioritize conservation actions using a habitat model able to explain abundance of Capercaillie at large scale. Pages poster in *Proceedings of XIII° International Symposium on Grouse*, Reykjavik, 3-13 sept. Abstract in *Grouse News* 51 : poster.

Gill, R. M. A., and R. J. Fuller. 2007. The effects of deer browsing on woodland structure and songbirds in lowland Britain. *Ibis* 149 : 119-127

Gjerde, I., and P. Wegge. 1989. Spacing pattern, habitat use and survival of Capercaillie in a fragmented winter habitat. *Ornis Scand.* 20:219-225.

GJERDE, I., P. WEGGE, O. PEDERSEN, and G. SOLBERG. 1984. Home range and habitat use of a local Capercaillie population during winter in S.E. Norway. Pages 247-260 in *Proceedings of 3rd Int.Grouse Symp.*:247-260.

Glad, A., B. Reineking, M. Montadert, A. Depraz, and J.-M. Monnet. 2019. Assessing the performance of object-oriented LiDAR predictors for forest bird habitat suitability modeling. *Remote Sensing in Ecology and Conservation*.

Graf, R., K. Bollmann, W. Suter, and H. Bugmann. 2005. The importance of spatial scale in habitat models: capercaillie in the Swiss Alps. *Landscape Ecology* 20:703-717.

Graf, R. F., K. Bollmann, H. Bugmann, and W. Suter. 2007. Forest and Landscape Structure as Predictors of Capercaillie Occurrence. *The Journal of Wildlife Management* 71:356-365.

Graf, R. F., S. Kramer-Schadt, N. Fernandez, and V. Grimm. 2007. What you see is where you go? Modeling dispersal in mountainous landscapes. *Landscape Ecology* 22:853-866.

Graf, R. F., W. Suter, and R. Hess. 2003. Le grand tétras-une espèce parapluie en voie de régression-mérite d'être sauvé. *La forêt* 2:10-12.

Hancock, M. 2010. Cattle, bilberry and capercaillie in a Scottish pinewood. *Grouse news* : p.10.

Hjeljord, O., P. Wegge, J. Rolstad, M. Ivanova, and A. B. Beshkarev. 2000. Spring-summer movements of male capercaillie *Tetrao urogallus*: A test of the 'landscape mosaic' hypothesis. *Wildlife Biology* 6 : 251-256.

Hurstel, A., and N. Braconnier. 2013. Impact des Ongulés sur la myrtille (*Vaccinium myrtillus*). Etude de cas sur le massif vosgien et influence sur la qualité des habitats du grand tétras (*Tetrao urogallus*). *Groupe tétras Vosges. Life Natura 2000*.

Kormann, U., K. Bollmann, F. Gugerli, and L. Excoffier. 2009. Avian dispersal in a mountain environment : landscape genetics in Capercaillie. Pages 145 in *Proceedings of 2nd european Congress of Conservation Biology: "Conservation Biology and Beyond: from science to practice*.145.

Kortmann et al. 2014. Effets du dépeuplement sur les habitats du Grand Tétras dans le Parc National de Bavière.

Kortland, K., and G. Findlay. 2010. Two capercaillie leks in novel habitat in Scotland. *Grouse news* 40:16-19.

- KURKI, S., A. NIKULA, P. HELLE, and H. LINDEN. 2000. LANDSCAPE FRAGMENTATION AND FOREST COMPOSITION EFFECTS ON GROUSE BREEDING SUCCESS IN BOREAL FORESTS. *Ecology* 81 : 1985-1997.
- LANDMANN, G. 1985. SYLVICULTURE ET GRAND TÉTRAS DANS LE MASSIF VOSGIEN : UN CONSTAT - DES PERSPECTIVES. *Revue forestière française* XXXVII : 135-152.
- LECLERCQ, B. 1987. Influence de quelques pratiques sylvicoles sur la qualité des biotopes à Grand tétras (*Tetrao urogallus*) dans le massif du Jura. *Acta Oecologica* vol n° 2, Oecol.génér.:237-246.
- Ludwig, G. X., R. V. Alatalo, P. Helle, K. Nissinen, and H. Siitari. 2008. Large-scale drainage and breeding success in boreal forest grouse. *Journal of Applied Ecology* 45:325-333.
- Ménoni, E., and J. Bougerol. 1993. Capercaillie population in forest fragmented by topography and human activities in the french Pyrenees. Pages (148-159) in Proceedings of XXI congress I.U.G.B. 2:(148-159).
- Ménoni, E., J. F. Brenot, and M. Catusse. (1994). Grand tétras et ski de fond. *Bulletin mensuel de l'ONC* 190 : 12-21.
- Ménoni, E., P. Landry, and C. Berducou. 1997. Habitat fragmentation and viability of capercaillie *Tetrao urogallus* populations in the french Pyrenees. *Wildlife Biology* 3:(277).
- Ménoni, E., N. Luigi, and F. Delfino. 2002. Is the capercaillie a relevant umbrella species in the french Pyrenees? in Proceedings of 9th international grouse symposium.
- Ménoni, E. & P. Defos du Rau. 2003. Démographie pyrénéenne du Grand Tétras.
- Ménoni, E., P. Defos du Rau, H. Brustel, A. Brin, L. Valladares, G. Coriol, L. Harvenc & J. L. Castel. (2005). Amélioration des habitats en faveur du Grand tétras et bénéfices escomptés sur la biodiversité. Faune sauvage. Hors-Série « Rapport Scientifique 2004 » : 65-68.
- Ménoni, E., D. Ilard, H. Verheyden, N. Morellet, L. Larrieu, E. Constantin, K. Saint-Hilaire, & D. Dubreuil. (2008). Cerf, troupeaux domestiques. Quels impacts sur l'habitat des Galliformes de montagne ? *Faune sauvage* 281 : 32-39.
- Ménoni, E. 2011. Habitat de la especie. in J. Canut, García-Ferré, D. &, andA. I., editors. *Manual de conservación y manejo del hábitat del urogallopirenaico*. Dirección General de Medio Natural y Política Forestal. Ministerio de Medio Ambiente, Medio Rural y Marino., Madrid.
- Ménoni, E., V. Favre-Ayala, R. Cantegrel, J. Revenga, J. Camprodón, D. Garcia, D. Campion, I. Afonso, & L. Riba. (2012). Réflexion technique pour la prise en compte du Grand tétras dans la gestion forestière pyrénéenne. Pau. FORESPiR, Union européenne, DREALMidi-Pyrénées.
- Ménoni, E. 2013. Handbook of technical considerations for management of Pyrenean capercaillie habitats. *Grouse news* 46:27-28.
- Miettinen, J. 2006. capercaillie as a tool of forest planning and as an indicator of forest environment quality. *Grouse news* 31 : 4-5.
- Mulhauser, B., and P. Junod. 2006. Sylviculture et revitalisation des habitats des tétraonidés dans le canton de Neuchâtel (Suisse). *Schweiz. Z. Forstwes.* 157 : 263-270.
- Office-National-des-Forêts. 1994. Gestion Forestière et Grand Tétras. Pyrénées. O.N.F. edition. O.N.F., Toulouse.
- Office National des Forêts. (2001). Reconstitution des forêts après tempêtes. Guide. ONF édition.

Pakkala, T., J. Pellikka, and H. Lindén. 2003. Capercaillie *Tetrao urogallus* - a good candidate for an umbrella species in taiga forests. *Wildlife Biology* 9:309-316.

Piedallu C., Dallery D., Bresson C., Legay M., Gégout, J.C., Pierrat R., 2022, Spatial vulnerability assessment of silver fir and Norway spruce dieback driven by climate warming, "Landscape Ecology" 38, 341-361

Piedallu C., Perez, V., Gégout J.C., Lebourgeois F., Bertrand R., 2009. Impact potentiel du changement climatique sur la distribution de l'Epicéa, du Sapin, du Hêtre, et du Chêne sessile en France. *Revue Forestière Française*, LXI - 6, 567-593.

Pollo, C. J., L. Robles, A. Garcia-Miranda, R. Otero, and J. R. Obeso. 2003. Variaciones en la densidad y asociaciones especiales entre ungulados silvestres y urogallo cantabrico. *Ecología* 17:199-206.

PETERHOFS, E., G. BAGRADE, G. DONE, M. LUKINS, A. ORNICANS, J. OZOLINŠ, D. PILATE, D.-E. RUNGIS, J. SUBA, and A. ZUNNA. 2019. SUITABILITY OF CAPERCAILLIE TETRAO UROGALLUS AS AN UMBRELLA SPECIES FOR LATVIAN BOREAL FORESTS. *in Proceedings of 34th IUGB Congress 2019*.

Quevedo, M., M. J. Banuelos, and J. R. Obeso. 2005. The decline of Cantabrian capercaillie: How much does habitat configuration matter? *Biological Conservation*. 127 : 190-200.

Quevedo, M., M. J. Banuelos, O. Saez, and J. R. Obeso. 2006. Habitat selection by Cantabrian capercaillie at the edge of species distribution. *Wildlife Biology* 12 : 267-276.

Rolstad, J. 1989. Habitat and range use of capercaillie *Tetrao urogallus* L. in southcentral Scandinavian boreal Forests, with special reference to the influence of modern forestry. Dr. agric. thesis, Agricuktural University of Norway, As.

Rolstad, J., I. Gjerde, V. S. Gunderson, and M. Saetersdal. 2002. Use of Indicator Species to Assess Forest Continuity: a critique. *Conservation Biology* 16 : 253-257.

Rolstad, J., E. Rolstad, and P. Wegge. 2007. capercaillie *Tetrao urogallus* lek formation in young forest. *Wildlife Biology* 13 : 59-67.

Rolstad, J., and P. Wegge. 1987a. Capercaillie habitat: a critical assessment of the role of old forest. Pages (242-254) *in Proceedings of 4th International Grouse Symposium (242-254)*.

Rolstad, J., and P. Wegge. 1987b. Distribution and size of capercaillie leks in relation to old forest fragmentation. *Oecologia (Berlin)* 72:(389-394).

Rolstad, J., and P. Wegge. 1989. Effects of logging on capercaillie (*Tetrao urogallus*) leks. IIIExtinction and recolonization of lek Populations in Relation to Clearfelling and Fragmentation of Old Forest. *Scandinavian journal on forest research* 4:129-135.

Rubiales, J. M., F. J. Ezquerra, F. Gómez-Manzaneque, S. G. Álvarez, I. García Amorena, and C. Morla. 2009. The long-term evolution of Cantabrian Mountain landscapes and its possible role in the capercaillie drama. *Grouse news* 38 : 9-11.

Sachot, S., N. Perrin, and C. Neet. 2003. Winter habitat selection by two sympatric forest grouse in western Switzerland: implications for conservation. *Biological Conservation* 112:373-382.

Sachot, S., and N. Perrin. 2004. Capercaillie (*Tetrao urogallus*) in western Switzerland. Viability and management of an endangered Grouse metapopulation. Pages 384-396 *in* U. Press, editor. *Species Conservation and Management-Case studies*. Akcakaya, H.R., Bergman, M.A. et al., New-York : Oxford.

Sachot, S., and C. Neet. 2006. Gestion sylvicole et Grand Tétras : les actions du canton de Vaud (Suisse). Nature Sciences Sociétés 14:60-62.

Sirkia, S., A. Lindén, P. Helle, A. Nikula, J. Knape, and H. Lindén. 2010. Are the declining trends in forest grouse populations due to changes in the forest age structure? A case study of Capercaille in Finland. Biological Conservation 143 : 1540-1548.

Sirkia, S., J. Pellikka, and H. Lindén. 2010. Balancing the needs of capercaillie (*Tetrao urogallus*) and moose (*Alces alces*) in large-scale human land use. Eur J Wildl Res 56 : 249-260.

Spidso, K., and H. Korsmo. 1994. Selection of feeding trees by capercaillie *Tetrao urogallus* in winter. Scandinavian Journal Forest researches 9 : 180-184.

Storch, I. 1993a. Habitat requirements of capercaillie. Pages (151-154) in Proceedings of Proceedings of the 6th International Grouse Symposium (151-154).

Storch, I. 1993b. Habitat selection by capercaillie in summer and autumn : is bilberry important ? Oecologia 95 : 257-265.

Storch, i. 1993c. Habitat Use and Spacing of Capercaille in relation to Forest Fragmentation patterns. PHD, Ludwig-Maximilian, Munich.

Storch, I. 1993d. Patterns and strategies of winter habitat selection in alpine capercaillie. *Ecography* 16 : 351-359.

STORCH, I. 1997. The importance of scale in habitat conservation for an endangered species: the capercaillie in central Europe. Pages 310- 330 in S. Verlag, editor. J. A. Bissonette, (ed). Wildlife and Landscape ecology: effects of pattern and scale. Springer Verlag, New York.

Suter, W., R. F. Graf, and R. Hess. 2002. Capercaillie (*Tetrao urogallus*) and avian biodiversity: testing the umbrella-species concept. *Conservation Biology* 16:778-788.

Swenson, J. E., and P. Angelstam. 1993. Habitat separation by sympatric forest grouse in fennoscandia in relation to boreal forest succession. Canadian journal of zoology. 71:(1303-1310).

Thiel, D., C. Unger, M. Kery, and L. Jenni. 2007b. Selection of night roosts in winter by capercaillie *Tetrao urogallus* in Central Europe. Wildlife Biology 13:73-86.

Viry, B., and C. Helderlé. 2012. Des forêts pour le Grand Tétras. Guide de sylviculture. Région Lorraine, GTV, CRPF, ONF, Life Tétras Vosges, AgroParisTech Interprint ed.

Wegge, P., M. H. Finne, and J. Rolstad. 2007. GPS satellite telemetry provides new insight into capercaillie *Tetrao urogallus* brood movements. Wildlife Biology 13:87-94.

WEGGE, P., I. GJERDE, L. KASTDALEN, J. ROLSTAD, and T. STORAAS. 1989. Does forest fragmentation increase the mortality rate of capercaillie? Pages 448-453 in Proceedings of 19th IUGB Congress. II : 448-453.

Wegge, P., and J. Rolstad. 1986. Size and spacing of capercaillie leks in relation to social behaviour and habitat. Behav Ecol Sociobiol 19 : 401-408.

Alimentation/nourriture

- Andreev, A. V. (1987). Ecological energetics of palearctic Tetraonidae in relation to chemical composition and digestibility of their winter diets. Canadian journal of zoology 66 : 1382-1388.
- Annila, E., N. P., and T. Levula.(2003). Observations on winter feeding of Capercaillie in fertilized Scots pine stands. In finnish, with english summary : Metson hakomisesta eri tavoin lannoitetuissa männiköissä. Suomen riista 49 : 25-31.
- Baines, D. (1991). Capercaillie decline and caterpillar abundance in highland pinewoods. Game conservancy annual review. 23 : 104-105.
- Baines, D., I. A. Wilson, and G. Beeley. (1996). Timing of breeding in Black grouse *Tetrao tetrix* and Capercaillie *Tetrao urogallus* and distribution of insect food for the chicks. *Ibis* 138 : 181-187.
- Borchtchevski, V. (2008). The May diet of Capercaillie (*Tetrao urogallus*) in an extensively logged area of NW Russia. *Ornis Fennica* 85 : 12 pp.
- Borchtchevski, V. (1994). Nourriture animale des juvéniles et des adultes Grands tétras (*Tetrao urogallus*) dans l'ouest de la taïga russe. *Gibier Faune Sauvage. Game and Wildlife*. 11(4) : 271-286.
- Borchtchevski, V. (2009). Space location of grouse birds as an adaptation to predator's pressure. Xle IGS, White Horse (Yukon).
- Borchtchevski, V. (2015). Does beginning of spring represent a critical period of annual cycle for Capercaillie (*Tetrao urogallus*, *Tetraonidae*, *Galliformes*) ? 94 (4): 455-465.
- Chua, P. Y.-S., Y. Lammers, E. Menoni, T. Ekrem, K. Bohmann, S. Boessenkool, and I. Greve-Alsos. 2021. Molecular dietary analyses of western capercaillies (*Tetrao urogallus*) reveal a diverse diet. *Environmental DNA* : 15 pp.
- Hissa, R., H. Rintamaeki, P. Virtanen, H. Linden, and V. Vihko. 1990. Energy reserves of the capercaillie. *Suomen riista* 36 : 62-71.
- Jacob, L. 1987. Le régime alimentaire du grand tétras (*Tetrao urogallus* L.). *Synthèse bibliographique. Gibier Faune Sauvage*. 4 : 429-448.
- Kastdalen, L. 1986. Food selection in capercaillie and black grouse chicks in South–East Norway., Oslo, Norway. Univeristy of Oslo.
- Moss, R., and I. Hanssen. 1980. Grouse nutrition. *Nutr Abstr Rev Ser B* 50 : 555-567.
- OBOTNIN, E. L. a. S. 2019. WILD BERRIES IN TETRAONIDAE FEEDING IN RUSSIAN TAIGA. Pages 150-151 in Proceedings of 34th IUGB Congress 2019. 150-151.
- Puissant, S. 2004. Les orthoptères comme outil d'aide à la gestion des sites de reproduction du grand tétras (massif du Canigou, Pyrénées-Orientales). *Matériaux Orthoptériques et Entomocénotiques* 9:(5-14).
- Rodriguez, A. E., and J. R. Obeso. 2000. Diet of the Cantabrian capercaillie: Geographic variation and energetic content. *Ardeola* 47:77-83.
- Saniga, M. 1998. Diet of the capercaillie (*Tetrao urogallus*) in a Central-European mixed spruce-beech-fir and mountain spruce forest. *Folia Zoologica* 47:115-124.

Selas, V. 2000. Population dynamics of Capercaillie in relation to bilberry production in southern Norway. *Wildlife Biology*, 6, 1-11.

Selas, V., et al. 2021. Bilberry seed production explains spatio temporal syncronicity in bank vole population fluctuations. *Ecological Research*, 36, 409-419.

Wegge, P., and L. KASTDALEN. 2008. Habitat and diet of young grouse broods: resource partitioning between Capercaillie (*Tetrao urogallus*) and Black Grouse (*Tetrao tetrix*) in boreal forests. *Journal of Ornithology* 149 : 237-244.

Wegge, P., T. Olstad, H. Gregersen, O. Hjeljord, and A. V. Sivkov. 2005b. Capercaillie broods in pristine boreal forest in northwestern Russia: the importance of insects and cover in habitat selection. *Canadian Journal of Zoology* 83:1547-1555.

Climat et changement climatique

Allstadt, A. J., A. M. Liebhold, D. M. Johnson, R. E. Davis, and K. J. Haynes. (2015). Temporal variations in the synchrony of weather and its consequences for spatiotemporal population dynamics. *Ecology* 96 : 2935-2946.

Archaud, F. (2008). Les oiseaux face au changement climatique. *Rendez-vous techniques* 20 : 65-68.

Badeau, V., J. L. Dupouey, C. Cluzeau, and J. Drapier. (2007). Aires potentielles de répartition des essences forestières d'ici 2100. *Rendez-Vous Techniques. Forêts et milieux naturels face aux changements climatiques* : 62-66.

Baines, D., N. J. Aebischer, and A. Macleod. (2016). Increased mammalian predators and climate change predict declines in breeding success and density of Capercaillie *Tetrao urogallus*, an old stand specialist, in fragmented Scottish forests. *Biodiversity Conservation* 25 : 2171-2186.

Barnagaud, J. Y., P. A. Crochet, Y. Magnani, A. Bernard Laurent, E. Ménoni, C. Novoa, and O. Gimenez. (2010). Short-term response to NAO but no long-term effects of climate change on the reproductive success of an alpine bird. *Journal of Ornithology* : 11 pp.

Beauvois, A. 2005. Influence des facteurs climatiques sur le succès de la reproduction chez *Tetrao urogallus* L. (Grand tétras) et *Lagopus mutus* L. (Lagopède alpin) dans les Pyrénées. Lyon, Université Claude Bernard et ONCFS : 31pp.

Borchtchevski, V. and V. A. Koshevoy. (2018). Does transforming geographical axes to climatic ones give new insights into the spatial distribution of the two Capercaillie species (*Tetrao urogallus*, *T. parvirostris*) ? *Grouse news* 55 : 17-24.

Borchtchevski, V. G., and V. A. Koshevoy. 2018. Does transforming geographical axes to climatic ones give new insights into the spatial distribution of the two capercaillie species (*Tetrao urogallus*, *T. parvirostris*)? *Grouse news* 55 : 17-24.

Braunisch, V., J. Coppes, R. Arlettaz, R. Suchant, H. Schmid, K. Bollmann. 2013. Selecting from correlated climate variables : a major source of uncertainty for predicting species distributions under climate change. *Ecography* 36 : 001 – 013.

Christensen, JH., et al. 2019. *Climate Dynamics*, 53 : 4857-4869.

Coppes J., Kammerle JL., Schroth KE., Braunish V., Suchant R. 2021. Weather conditions explain reproductive success and advancement of the breeding season in Western Capercaillie (*Tetrao urogallus*). *Ibis* : 163, 990–1003.

Crick, H. Q. P. 2004. The impact of climate change on birds. *Ibis* 146:48–56. DOI 10.1007/s10144-010-0255-0.

FLETCHER, K., D. HOWARTH, A. KIRBY, R. DUNN, and A. SMITH. 2013. Effect of climate change on breeding phenology, clutch size and chick survival of an upland bird. *Ibis* 155:456–463.

Fuller, R. J. 2001. Responses of woodland birds to increasing numbers of deer: a review of evidence and mechanisms. *Forestry* 74:(289-298).

González, M. A. 2014. Biology and conservation of the capercaillie (*Tetrao urogallus*) in a Mediterranean environment. Thesis summary. *Grouse news* 47:30.

Gonzalez, M. A., P. P. Olea, L. Robles, and V. Ena. 2010. The Mediterranean Quercus pyrenaica oak forest: a new habitat for the Capercaillie? *Journal of Ornithology* DOI 10.1007/s10336-010-0528-6:6 pp.

Hancock, M., 2009, Capercaillie habitat benefits from an unusual climatic event in Scotland. *Grouse news* 37 : 7-8.

Heljord, et Loe, L-E. 2022. The role of climate and alternative prey in explaining 142 years of declining Willow grouse hunting bag.

Howard, C., Marjakangas, E. L., Morán-Ordóñez, A., Milanesi, P., Abuladze, A., Aghababyan, K., ... & Willis, S. G. (2023). Local colonisations and extinctions of European birds are poorly explained by changes in climate suitability. *Nature Communications*, 14(1), 4304.

Kvasnes, M., et al. 2014. Large scale climate vulnerability and rodent abundance modulates recruitment rates in Willow Ptarmigan (*lagopus lagopus*). *Jour. of Ornithology*, 155, 891-903.

Leclercq, B. 1984. Le climat et la régression du grand tétras. *Bulletin Mensuel de l'ONC* 79:15-21.

Leclercq, B., and E. Ménoni. 2019. Effets des changements climatiques sur l'écologie de quelques tétraonidés. Bourgogne-Franche-Comté Nature Actes des 15°Rencontres Bourgogne-Franche-Comté Nature : 168-186.

Le Gallic, J. 2021. Phénologie de la reproduction du Grand Tétras dans les Vosges. Stage de Master EEE. Groupe Tétras Vosges.

Ludwig, G. X. 2009. Impact of climatic and environmental change on Finnish grouse populations. *Suomen riista* 55:7-18.

Ludwig, G. X., R. V. Alatalo, P. Helle, H. Linden, J. Lindstrom, and S. H. 2006. Short- and long-term population dynamical consequences of asymmetric climate change in black grouse. *Proceeding of the Royal Society Bulletin* 273 : 2009-2016.

Ménoni, E. & C. Novoa (2007). Les effets des changements climatiques sur les oiseaux : l'exemple des Galliformes de montagne en France. *Rendez- Vous Techniques. Forêts et milieux naturels face aux changements climatiques*. Hors-série n°3 : 53-61.

Ménoni, E. 2013. Les grands tétras s'accouplent de plus en plus tôt : un effet mesurable du changement climatique. Rapport Scientifique de l'oncfs 2013 : p.43.

Ménoni, E., M. Montadert, B. Leclercq, A. Hurstel, & K. Dillet. (2012). Change in mating and breeding time of the Capercaillie in France, in relation to the change of the phenology of spring vegetation. in Proceedings of XII International Symposium on Grouse. Abstract.

Menoni, E., and C. Novoa. 2020. Effets du changement climatique sur les espèces animales dites « arctico-alpine », exemple des espèces emblématiques grand tétras et lagopède alpin. in Proceedings of Carrefour du climat.

Moss, R. 1985. Rain, breeding success and distribution of Capercaillie *Tetrao urogallus* and Black Grouse *tetrix* in Scotland. *Ibis* 128 : 65-72.

Niemelä, T. E., and H. Lindén. 2009. Capercaillie and the changing winter climate. *Suomen riista* 55:19-34.

Novoa, C., G. Astruc, J.-F. Desmet, and A. Besnard. 2016. No short-term effects of climate change on the breeding of Rock Ptarmigan in the French Alps and Pyrenees. *Journal of Ornithology* 157 : 797-810.

Roy S. 2020. Évolution climatique pour la Région Grand Est : plusieurs scénarios. Météo France.
<https://www.grandest.fr/wp-content/uploads/2020/02/roy-sophie.pdf>

Selas, V., G. A. Sonerud, E. Framstad, J. A. Kalas, S. Kobro, H. B. Pedersen, T. K. Spidsø, and Ø. Wiig. 2011. Climate change in Norway : warm summers limit grouse reproduction. *Popul Ecol* 53 : 361–371.

Sitzia, T., T. Campagnaro, M. Dainese, F. Dallabrida, S. M. Mattioli, and A. Portaccio. 2019. Altitudinal Shift of *Tetrao urogallus* in an Alpine Natura 2000 Site: Implications for Habitat Restoration. *Applied Science* 9, 1164.

Sorensen-Forbey, J., N. L. Wiggins, G. G. Frye, and J. W. Connelly. 2013. Hungry grouse in a warming world: emerging risks from plant chemical defenses and climate change. *Wildlife Biology* 19:374-381.

Swenson, J. et al., 1994. Effects on Hazel Grouse reproduction : an allometric perspective. *J. of Avian Biology*, 25 : 1-8.

Wegge, P., T. Vesterås, and J. Rolstad. 2010. Is the timing of breeding and subsequent hatching in boreal forest grouse adapted to insect food for the chicks? *Ornis Fennica* 10 press:21 pp.

Wegge, P. 2017. Climate change and bird reproduction: warmer springs benefit breeding success in boreal forest grouse. *Proc. R. Soc. B* 284: 15-28.

Zbinden N., Korner-Nievergelt F., Tettamanti F., Keller V. 2022. Long-term trends of reproductive success of black grouse *Lyrurus tetrix* in the southern Swiss Alps in relation to changes in climate and habitat. *Wildlife Biology*, e01022.

Derangement /perturbation

Arlettaz, R., P. Patthey, M. Baltic, T. Leu, M. Schaub, R. Palme, and S. Jenni-Eiermann. 2006. Spreading free-riding snow sports represent a novel serious threat for wildlife. Proceeding of the Royal Society B. www.journals.royalsoc.ac.uk, published online.

Arlettaz, R., P. Patthey, and V. Braunisch. 2012. Impacts of Outdoor Winter Recreation on Alpine Wildlife and Mitigation Approaches: A Case Study of the Black Grouse. Pages 137-154 in *The Impact of Skiing on Mountain Environments*. Bentham Science Publishers.

Baines, D. (2003). Does increased access damage success of nesting birds ? The Game Conservancy Trust (34) : 31-34.

Baines, D., M. Richardson. 2007. "An experiential assessment of the potential effects of human disturbance on Black Grouse in the North Pennines ». Ibis 149, suppl 1.

Bal, G., L. Bacon, E. Menoni ; C. Calenge, A. Million, A Besnard. 2021. Modélisation de la dynamique du Grand Tétras des Pyrénées françaises pour sa gestion adaptative. Technical report MNHN et OFB. Rendu public en 2021, in press.

Baltic et al. 2005. "A non invasive technique to evaluate human generated stress in Black Grouse Ann. N.Y. Acad.Science, doi : 10.1196/annals. 1343. 008.

Belleau, E. (2006). Parasitology as a contribution towards measuring human-generated stress in black grouse." Grouse news 32 : 11-12.

Blanc, R., M. Guillemain, J. B. Mouronval, D. Desmonts, and H. Fritz. (2006). Effects of non-consumptive leisure disturbance to wildlife. Revue d'écologie (Terre et Vie) 61 : 117-133.

Bortchevski et al., 2003, Does fragmentation by logging reduce grouse reproductive success in boreal forests ? Wildlife Biology, 9, 275-283.

Brenot, J. F., and E. Ménoni. 1999. RESPONSE OF A CAPERCAILLIE POPULATION IN RELATION TO CONSTRUCTION OF A SKI STATION IN THE PYRENEES. Pages 1p. in Proceedings of The 8th International Grouse Symposium.1p.

Brenot, J. F., M. Catusse, and E. Ménoni. 1996. Effets de la station de ski de fond du plateau de Beille (Ariège) sur une importante population de grand tétras (*Tetrao urogallus*). Alauda 64:(249-260).

Chesnais M., 2023, Suivi de la fréquentation sur la Réserve Naturelle Nationale du Tanet-Gazon du Faing, Conservatoire d'espaces naturels de Lorraine. Document de travail en cours de redaction.

Coppes, J., and V. Braunisch. 2013. Managing visitors in nature areas: where do they leave the trails? A spatial model. Wildlife Biology 19:1-11.

Coppes, J., J. Ehrlacher, R. Suchant, and V. Braunisch. 2017. Outdoor recreation causes effective habitat reduction in Capercaillie *Tetrao urogallus*: a major threat for geographically restricted populations. Journal of Avian Biology:41.

Coppes, J., J. L. Kämmerle, M. Willert, A. Kohnen, R. Palme, and V. Braunisch. 2018. The importance of individual heterogeneity for interpreting faecal glucocorticoid metabolite levels in wildlife studies. Grouse news : 25.

LABIGAND, G., M. MONTADERT, C. GUILLEMOT, R. SUCHANT, H. WEISS, F. PREISS, and J. POIROT. 1995. Etude comparative Forêt Noire- Vosges- Jura. Pages 88- 107 in Proceedings of Le grand tétras question de survie.:88- 107.

LABIGAND, G., and M. MUNIER. 1989. Grand tétras et tourisme hivernal, historique d'une place de chant dans les Hautes Vosges. Ciconia 13 : 19- 31.

LECLERCQ, B. 1985. Influence des routes et voies de pénétration humaine sur les comportements de grand tétras et de gelinottes dans le haut Jura français. Pages 197-203 in Proceedings of Colloque "routes et faune sauvage ".197-203.

Lingg. 2008. Is wildlife inhabiting snow sport areas less fear than in undisturbed areas ? A case of study of the alpine Black Grouse. Bern University.

Ménoni, E., P. Defos du Rau, & P. Blanc. (2007). Extension des domaines skiables et Grand tétras : l'expertise ONCFS. Rapport scientifique de l'ONCFS : 10-16.

Ménoni, E. & J. Fortin. (2015). Emprise des loisirs de plein air sur les habitats des Galliformes de montagne dans les Pyrénées françaises. Faune Sauvage 306 : 38-43.

Ménoni, E. & Y. Magnani. (1998). Human disturbance of grouse in France. Grouse news 15 : 4-8.

Montadert, M. 2007. " Le dérangement : définition et approche scientifique développées pour évaluer son impact". Faune Sauvage n° ?, p ?.

Montadert, M. 2013. Tétras-lyre et dérangement touristique : Synthèse bibliographique. Fédération Départementale des Chasseurs de l'Isère. 44 pp.

Watson, A., and R. Moss. 2004. Impacts of ski development on Ptarmigan (*lagopus mutus*) at Cairn Gorm Scotland. Biological Conservation, 116. 267-275

Moss, R., F. Leckie, A. Biggins, T. Poole, D. Baines, and K. Kortland. 2014. Impacts of human disturbance on Capercaillie *Tetrao urogallus* distribution and demography in Scottish woodland. Wildlife Biology 20:1-18.

Parc naturel régional des Ballons des Vosges. 2021. Suivi de la fréquentation par appareils photos à déclenchement automatique sur deux sites de présence du Grand Tétras, 2020-2021.

Poirot, J. 2023. Suivi d'une place de chant de Grand tétras par appareils photos à déclenchement automatique en lien avec les dé/confinements. Tétrarchives.

Poirot, J. 2022. Suivi de la grande faune par deux appareils photos à déclenchement automatique, en lien avec la fréquentation humaine. Tétrarchives.

Porkert, J. 1978. The influence of human factors on tetraonid populations in Czechoslovakia. Pages 74-82. in Proceedings of First International Grouse Symposium.74-82.

Rupf, R., M. Wytttenbach, D. Köchli, M. Hediger, S. Lauber, P. Ochsner, and R. Graf. 2011. Assessing the spatio-temporal pattern of winter sports activities to minimize disturbance in capercaillie habitats. e c o.mo n t 3.

Storch, I. 2013. Human disturbance of grouse - why and when? Wildlife Biology 19:390-403.

Summers, R. W., J. McFarlane, and J. W. Pearce-Higgins. 2007. Measuring avoidance by capercaillies *Tetrao urogallus* of woodland close to tracks. Wildlife Biology 13:19-27.

Thiel, D., E. Ménoni, J. F. Brenot, and L. Jenni. 2007. Effects of recreation and hunting on flushing distance of capercaillie. *Journal of Wildlife Management* 71:1784-1792.

Thiel, D. 2007a. Behavioral and Physiological Effects in Capercaillie (*Tetrao urogallus*) Caused by Human Disturbance., Zurich.

Thiel, D. 2007b. Measuring Corticosterone Metabolites in Droppings of Capercaillies (*Tetrao urogallus*). *Annals of the New York Academy of Sciences* 1046:96-108.

THIEL, D., S. JENNI-EIERMANN, R. PALME, and L. JENNI. 2011. Winter tourism increases stress hormone levels in the Capercaillie *Tetrao urogallus*. *Ibis* 153:122-133.

Thiel, D., S. Jenni-Eirmann, V. Braunisch, R. Palme, and L. Jenni. 2008. Ski tourism affects habitat use and evokes physiological stress response in Capercaillie *Tetrao urogallus*: a new methodological approach. *Journal of Applied Ecology* 45:845-853.

Wegge, P. and Rolstad J., 2011, Clearcutting and eurasian boreal grouse : Long term monitoring reveals unexpected effects on their population performances. *Forest Management*, 261.

Zeitler, A., 2007, Habituation of Black Grouse to humans in the Bavarian Alps ? 4th International Black Grouse Conference, Vienne, 2007.

Zeitler, A., 2000, Human disturbances, behaviour and spatial distribution of Black Grouse in skiing areas in the Bavarian Alps. *Cahier d'Ethologie*, 20, 381-402.

Mortalité dues aux infrastructures (câbles, clôtures, éoliennes)

Baines, D. and M. Andrew (2003). Marking of deer fences to reduce frequency of collisions by woodland Grouse. *Biological Conservation* 110 : 169-176.

Baines, D. and R. W. Summers (1997). Assessment of bird collisions with deer fences in Scottish forests. *Journal of applied ecology*. 34 : 941-948.

Bech, N., S. Beltran, J. Boissier, J. F. Allienne, J. Resseguyer, and C. Novoa. (2012). Bird mortality related to collisions with ski –lift cables : do we estimate just the tip of the iceberg ? *Animal Biodiversity and Conservation* 35 : 95-98.

Bevanger, K. (1988). Transmission line wirestrikes of Capercaillie and Black grouse in central Norwegian coniferous forests. *Okoforsk rapp*. 9: 1-53.

Bevanger, K. (1995). Estimates and population consequences of Tetraonid mortality caused by collisions with high tension power lines in Norway. *Journal of applied ecology* 32 : 745-753.

Bevanger, K. and H. Broseth (2000). Bird collisions with power lines-an experiment with ptarmigan (*Lagopus spp.*). *Biological Conservation* 99 : 341-346.

Buffet, N., and E. Dumont Dayot. 2013. Bird Collisions with Overhead Ski-Cables: A Reducible Source of Mortality. Pages 123-136 in *The Impacts of Skiing on Mountain Environments*. Rixen Christian and Rolando Antonio (Eds).

Christiansen, T. 2009. Fence Marking to Reduce Greater Sage-grouse (*Centrocercus urophasianus*) Collisions and Mortality near Farson, Wyoming – Summary of Interim Results. University of Oklahoma's Sutton Avian Research Center.

Dutry, M. 2020. Etude pour la reduction de certains facteurs limitants à la conservation du Grand Tétras dans le massif des Vosges. Mémoire de fins d'études. PNRBV. Master 2 Biodiversité, Ecologie, Evolution ; Université de Perpignan, Via dolomitia.

González, M. A. 2018. Female Cantabrian capercaillie dead by collision with wind turbine. Grouse news 55:15-17.

Tetrao urogallus ; quel impact de la chasse et des infrastructures ? Premières rencontres naturalistes de Midi-Pyrénées.Cahors.

Novoa, C., E. Hansen, & E. Ménoni. (1990). La mortalité de trois espèces de Galliformes par collision dans les câbles ; résultats d'une enquête pyrénéenne. Bulletin mensuel de l'ONC 151 : 17- 22.

Peny, S. J. 1995. Assessment of fence collisions by grouse species in Scotland. Research information of the Research Division of the Forestry Commission. 264:(1-9).

Stevens, B. S. 2011. Impacts of fences on Greater Sage-Grouse in Idaho: collision, mitigation, and spatial ecology. M. Sc. Thesis. University of Idaho. Grouse news 44.

Warren, P., and D. Baines. 2002. Dispersal, survival and causes of mortality in black grouse Tetrao tetrix in northern England. Wildlife Biology 8:(129-135).

Wolfe, D., M. A. Patten, and S. K. Sherrod. 2009. Reducing Grouse Collision Mortality by marking Fences (Oklahoma). Ecological restoration 27:141-143.

Wolfe, D. H., M. A. Patten, E. Shochat, C. L. Pruett, and S. K. Sherrod. 2007. Causes and patterns of mortality in lesser prairie-chickens *Tympanuchus pallidicinctus* and implications for management. Wildlife Biology 13:95-104.

Génétique

Alda, F., P. Sastre, P. J. De La Cruz-Cardiel, and I. Doadrio. (2010). Population genetics of the endangered Cantabrian Capercaillie in northern Spain. Animal Conservation. Print ISSN 1367-9430.

Alda, F., M. A. González, P. P. Olea, V. Ena, R. Godinho, and S. V. Drovetski. (2013). Genetic diversity, structure and conservation of the endangered Cantabrian Capercaillie in a unique peripheral habitat. Eur J Wildl Res, 59 : 719-728.

ALEIX-MATA, G., F. J. RUIZ-RUANO, J. M. PÉREZ, M. SARASA, and A. SÁNCHEZ. 2019. Complete mitochondrial genome of the Western Capercaillie *Tetrao urogallus* (Phasianidae, Tetraoninae). Zootaxa 4550 : 585-593.

Bajc, M., M. Cas, D. Ballian, S. Kunovac, G. Zubic, M. Grubec, P. Zhelev, L. Paule, T. Grebenc, and H. Kraigher. (2011). Genetic differentiation of the western Capercaillie highlights the importance of South-Eastern Europe for understanding the species phylogeography. PlosOne 6.

Blondel, J. (2005). De l'histoire naturelle à la génétique de la gestion à la conservation. Faune sauvage 265 (Mars 2005) : 1-2.

Borchtchevski, V. G., and I. A. Khomyakova. 2018. THE capercaillie (*Tetrao urogallus*) bod y weight: spata variability in the west of the range. УДК 598.261.6, 591.134.5.

Bouzat, J. L., H. H. Cheng, H. A. Lewin, R. L. Westemeier, J. D. Brawn, and K. N. Paige. (1998). Genetic evaluation of a demographic bottleneck in the greater prairie chicken. *Conservation Biology* 12 : 836-843.

Bouzat, J. L., J. A. Johnson, J. E. Toepfer, S. A. Simpson, T. L. Esker, and R. L. Westemeier. (2009). Beyond the beneficial effects of translocations as an effective tool for the genetic restoration of isolated populations. *Conservation Genetic* 10 : 191-201.

Charge, R., C. Teplitsky, Y. Hingrat, M. Saint Jalme, F. Lacroix, and G. Sorci. 2013. Quantitative genetics of sexual display, ejaculate quality and size in a lekking species. *Journal of Animal Ecology* 82 : 399-407.

Cayuela, H., L. Boualit, M. Laporte, J. G. Prunier, F. Preiss, A. Laurent, F. Foletti, J. Clobert, and G. Jacob. 2019a. Kin dependent dispersal influences relatedness and genetic structuring in a lek system. *Oecologia* : 17pp.

Cayuela et al. 2021. Demography genetics and decline of a spatially structured population of a lekking bird. *Oecologia* 195, 117-129.

Duriez, O., J. M. Sachet, E. Ménoni, N. Pidancier, C. Miquel, and P. Taberlet. 2007. Phylogeography of the capercaillie in Eurasia: what is the conservation status in the Pyrenees and Cantabrian Mounts? *Conservation Genetic* DOI 10-1007/s10592-006-9165-2 : 14 pp.

Escoda, L., J. Castresana. 2021. Analisis genomico de las poblaciones europeas de urogallo y evaluacion del estado de conservacion desde el punto de vista genetico de las poblaciones amenazadas en Espana. Instituto de biología evolutiva, Barcelona.

Foletti, F. 2014. Non-invading genetic monitoring of capercaillie in the wild : individual tracking and breeding success., Master thesis-University of Fribourg, Domain of Ecology and Evolution, Department of Biology.

Foletti, F., and G. Jacob. 2016. Étude génétique de la population de Grand tétras (*Tetrao urogallus*) dans le Parc National des Cévennes. Université de Fribourg, ONCFS, Parc National des Cévennes. 28 pp.

Groupe Tétras Vosges. 2020. Estimer le nombre de mâles de Grand Tétras : comptage au chant ou génotypage ? Brèves du Groupe Tétras Vosges : 1 p.

Hurstel, A., and G. Jacob. 2014. Étude génétique de la population de Grand tétras (*Tetrao urogallus major*) des Vosges.

Hammerly J.-L. et al. 2013. A comparison of pedigree and DNA based measures for identifying inbreeding depression in the endangered. *Attawer's Prairie chicken Molecular Ecology*. 22 5313-5328.

Isomursu M. et al. 2012. Susceptibility to intestinal parasites and juvenile survival are correlated with multilocus microsatellites heterozygosity in the Capercaillie. *Ornis Fennica*, Janv 2012.

Jacob G., F. Foletti. 2022. Grand Tétras et génétique : apport de la génétique pour l'étude et la gestion des populations menacées. L'érosion de la diversité génétique menace la survie de la population jurassienne du Grand Tétras. Présentation au Comité Technique du PNA Grand Tétras déclinaison Jura.

Jacob G., F. Foletti. 2022. Mitteleuropäisches Raufussuhner Expertentreffen. Bad Peterstal.

- Karl Larson, J. 2006. Population fragmentation and genetic variation in grouse. Grouse news 31 : p 29.
- Larson, K. 2006. Population fragmentation and genetic variation in grouse. Grouse news 31 : p.29.
- Liukkonen-Antilla, T., O. Rätti, L. Kvist, and H. P. 2004. Lack of structuring and subspecies differentiation in the capercaillie (*Tetrao urogallus*) in Finland. Ann. Zool. Fennici 41 : 619-633.
- Lucchini, V., J. Hoglund, S. Klaus, J. Swenson, and E. Randi. 2001. Historical biogeography and a mitochondrial DNA phylogeny of grouse and ptarmigan. Molecular Phylogenetics and Evolution 20:149-162.
- Massard, P. 2017. Premiers résultats de six années de suivi génétique de la population de Grand Tétras (*Tetrao urogallus major*) sur le massif vosgien. Stage de Master. Groupe Tétras Vosges.
- Novoa, C., N. Bech, J. Resseguiier, R. Martinez-Vidal, D. Garcia-Ferré, J. Sola-de-la-Torre, and J. Boissier. 2014. A translocation experiment for improving the genetic diversity of an isolated population of Pyrenean rock ptarmigan (*Lagopus muta pyrenaica*). Grouse news 47 : 11-18.
- Regnault, S., and E. Ménoni. 2005. Génétique des populations Jurassiennes et Pyrénéennes de Grand tétras. Faune sauvage 265, spécial génétique : p.31.
- Regnaut, S. 2004. Population genetics of Capercaillie (*Tetrao urogallus*) in the Jura and the Pyrenees: a non-invasive approach to avian conservation genetics. Thèse de Doctorat ès sciences de la vie (PhD), Faculté de Biologie et de Médecine., Lausanne.
- Regnaut, S., P. Christe, M. Chapuisat, and L. Fumagalli. 2006. Genotyping faeces reveals facultative kin association on capercaillie's leks. Conservation genetics 7 : 665-674.
- Rodriguez-Munoz, R., P. M. Mirol, G. Segelbacher, A. Fernandez, and T. Tregenza. 2006. Genetic differentiation of an endangered capercaillie (*Tetrao urogallus*) population at the Southern edge of the species range. Conservation Genetic DOI 10 : 12 pp.
- Segelbacher, G., J. Hoglund, and I. Storch. 2003. From connectivity to isolation: genetic consequences of population fragmentation in capercaillie across Europe. Molecular Ecology 10:8 pp.
- SEGELBACHER, G., S. MANEL, and J. TOMIUK. 2008. Temporal and spatial analyses disclose consequences of habitat fragmentation on the genetic diversity in capercaillie (*Tetrao urogallus*). Molecular Ecology 17 : 2356-2367.
- Segelbacher, G., and S. Piertney. 2007. Phylogeography of the European capercaillie (*Tetrao urogallus*) and its implication for conservation. Journal of Ornithology 148 (Suppl 2) : S269-S274.
- Segelbacher, G., P. Wegge, A. V. Sivkov, and J. Höglund. 2007. Kin groups in closely spaced capercaillie leks. Journal of Ornithology 148 : 79-84.
- Storch, I., and G. Segelbacher. 2000. Genetic correlates of spatial population structure in central European capercaillie *Tetrao urogallus* and black grouse *T. tetrix*: a project in progress. Wildlife Biology 6: 305-310.
- Westemeier et al. 1998. Tracking the long term decline and recovery of an isolated population (prairie chicken). Science (NY).

Reintroduction /élevage

Angelstam, P. and F. Sandegren (1981). A release of penreared Capercaillie in central Sweden- survival, dispersal and choice of habitat. 2nd international symposium on grouse, Edinburgh, WorldPheasant Association.

Baydack, R. and B. Vodehnal (2005). A management strategy for North American grouse. Grouse news 30 : 9-10.

Buchanan Stevenson, G. 2007. An historical account of the social and ecological causes of capercaillies (*Tetrao urogallus*) extinction and reintroduction in Scotland. Thesis submitted for the degree of Doctor of Philosophy at the University of Stirling.

Gigot, G., A. Andrieu, J. Comolet-Tirman, S. Wroza. 2023. Synthèse sur les scenarios de renforcement de Grand tétras en France. Rapport Patri-Nat.

Kowalczyk, A., E. Lukaszewicz, and Z. Rzonca. 2012. New, Simply and effective freezing method of capercaillie (*Tetrao urogallus L.*) semen. Pages 1 in Proceedings of 12th International Symposium on Grouse. 1.

Lukaszewicz, E., A. Kowalczyk, and Z. Rzońca. 2011. Successful semen collection from Capercaillie (*Tetrao urogallus L.*) kept in an aviary system. *Ornis Fennica* 88 : 110-115.

Ménoni, E., K. Foulché. 2021. Faisabilité, modalités pratiques et mesures d'accompagnement d'une translocation de Grand tétras dans les Vosges. DRAS-OFB.

Merta, D., J. Kobielski, J. Theuerkauf, and R. Gula. 2016. Towards a successful reintroduction of capercaillies – activity, movements and diet of young released to the Lower Silesia Forest, Poland. *Wildlife Biology* 22 : 130-135.

Nappée, C., and G. Douhéret. 2004. Development of the reintroduced Capercaillie population in the Parc National des Cévennes. Grouse news 28 : 9-11.

Nappée, C. 2008. Le grand tétras (*Tetrao urogallus*) dans les Cévennes : histoire d'une réintroduction difficile. *Ornithos* 15 : 282-293.

Nappée, C. 2009. The reintroduced population of capercaillie in the Parc National des Cévennes is still alive! Grouse news 38 : 29-30.

ONCFS. 2017. Note sur les captures de galliformes de montagne : marche à suivre de la capture au relâcher de l'oiseau.

OFB, 2023, Contribution française au programme de sauvegarde du Grand Tétras des Monts Cantabriques, fiche de communication et journée technique.

Parc naturel régional des Ballons des Vosges. 2023. Retours d'expérience des voyages d'études à la préparation d'un projet de renforcement de Grand Tétras dans les Vosges.

Quevedo, M., R. Rodríguez-Muñoz, M. Banuelos, and A. Fernández-Gil. 2005. A captive breeding programme for Cantabrian capercaillie: does it make any sense? Grouse news 30:10-13.

Romanov, A. N. 1988. Capercaillie. Agropromizdat, Moscow edition.

Rzońca, Z., E. Lukaszewicz, and A. Kowalczyk. 2012. Protection, reproduction and reintroduction of capercaillie in the Forestry Wiśla, Poland. Grouse news 43:17-20.

Sedon, P. J., et al. 2014. Reintroducing resurrected species : selecting DeExtinction candidates. Trends in Ecology and Evolution. March, 2014, vol. 29, no.3.

Seiler, Ch., P. Angelstam, H.-H. Bergmann. 2000. Conservation releases of captive-reared grouse in Europe : what do we known and what do we need ? Cahiers d'éthologie, 20 (2-3-4) : 235 – 252.

Siano, R. 2005. The re-introduction of capercaillie (*Tetrao urogallus* L.) in the Harz Mountains National Park – state of the art and a PhD project in process. Grouse news 29:16-18.

Sokos, C. K., P. K. Birtsas, and E. P. Tsachalidis. 2008. The aims of galliforms release and choice of techniques. Wildlife Biology 14:412-422.

Snyder et al., 1999. Translocation Histories of Prairie Grouse in the United States.

IUCN. Guidelines for reintroductions and other conservation translocations. Species survival commission.

WARREN, P., P. J. C. WHITE, D. BAINES, F. ATTERTON, and M. J. BROWN. 2013. Variations in Black Grouse *Tetrao tetrix* winter survival in a year with prolonged snow cover. Bird Study 60 : 257-263.

Warren, P. et al. 2017. Expanding the range of black grouse *Tetrao tetrix* in northern England through translocating wild males. Wildlife Biology 2017 : wlb.00242.

World-Pheasant-Association, and IUCN/SSC-Re-introduction-Specialist-Group. 2009. Guidelines for the Re-introduction of Galliformes for Conservation Purposes. Gland, Switzerland: IUCN and Newcastle-upon-Tyne, UK: World Pheasant Association. World Pheasant Association and IUCN/SSC Re-introduction Specialist Group edition.

Zimmermann, B., and L. Thielemann. 2018. Results of a pilot study on the reintroduction of Capercaillie *Tetrao urogallus* to Brandenburg, Germany. in German with English summary. VOGELWELT 138 : 29 - 53.